

Post-Operative Instructions for Knee Arthroscopy

Dr. Bushnell

□ Meniscectomy,	
Progressively increase weight-bearing while us	ing the crutches
Use crutches until you can walk without a limp	and without significant pain
While sitting, straighten and bend the knee sev	eral times per hour
You can shower on the 3rd day after surgery, do	not submerge extremity for 2 weeks
Steri-strips can be removed 5 days after surger	γ
Your sutures are under the skin and will dissolv	e on their own
Do not put Neosporin or other ointment on the	incision sites more than once daily
Return to daily activity as your are comfortable	
☐ ACL, meniscectomy, meniscus repa	ir, microfracture, MPFL reconstruction
Knee brace should be used at all times	
Use crutches and put	weight through the operative leg
You can shower on the 3 rd day after surgery, do	not submerge extremity for 2 weeks
Steri-strips can be removed 5 days after surger	У
Your sutures are under the skin and will dissolv	re on their own
Do not put Neosporin or other ointment on the	incision sites more than once daily
Begin Physical Therapy	
-Kinex Equipment	
You can use the TENS unit and cold pack as of Begin CPM and pro	

DVT/blood clot precaution

Use compression stockings for 2 weeks
Take 325mg(adult strength) aspirin 2 times per day for 2 weeks

Pain Medications

Call the clinic (706-236-6362) during office hours to request pain medication refills Medication refills may take up to 24 hours to be approved Per clinic policy- NO medication refills will be made after hours or on weekends